

**Original article:**

## **Evaluation of Platelet Indices among Patients with Dengue Infection attending OBGY OPD/IPD in District Hospital , Churu , Rajasthan**

**Dr Gogaram<sup>1#</sup>, Dr Hanuman Prasad<sup>2\*</sup>, Dr Sarad Misha<sup>1</sup>**

#: First author; \* : Corresponding author

1: Principal specialist, Obs &Gyn, DBH Churu; 2: Associate professor, Churu Medical College.

---

### **Abstract**

**Introduction:** Dengue is a global endemic and most prevalent human arbovirus diseases, occurring in tropical and subtropical countries of the world where over 2.5 billion people are at risk of infection. During the past few years, the characteristics of dengue in India appear to have changed. One of the most common laboratory findings in dengue is thrombocytopenia. The complex mechanism of thrombocytopenia remains unclear. Recently, novel platelet indices such as MPV, PDW, and P-LCR have been investigated as prospective platelet activation markers.

**Objectives:** The aim of our study was to investigate the platelet indices in patients compared to the control and to assess the role of it in the severity dengue infection.

**Materials and methods:** This study was conducted prospectively for a period from July 2015 to November 2015 during the recent outbreak of dengue. The platelet parameters were measured by the semi automated hematology analyzer on venous samples collected in EDTA from 200 patients as a clinical sample and 200 apparent healthy normal individuals as control sample. The inclusion criteria were all patients with clinical features and serologically positive dengue infection included. The exclusion criteria include patient's serologically negative dengue and if routine laboratory testing suggested a bacterial, parasite or any viral infection other than dengue infection or any other disease.

**Results:** The differences between the patient group and the control group in MPV, PCT and PLT were found to be significantly lower in the patient group ( $P < 0.0001$ ), while the differences between patient group and control in PDW and P-LCR were found to be significant higher in patient group ( $P < 0.0001$ ). There was a significantly lower Platelet count and PCT in DHF group as compared to DF group ( $P < 0.0016$  and  $< 0.0166$  respectively) and significantly higher PDW in DHF group as compared DF group ( $P < 0.0008$ ). There was a significant negative correlation with mean platelet count and PDW and P-LCR and significant positive correlation with mean platelet count and PCT. High P-LCR and low PCT are more sensitive for DHF than DF and equally specific for both DF and DHF. High PDW is more sensitive for DHF and more specific for DF and low MPV is more sensitive for DF and equally specific for DF and DHF.

**Conclusion:** In conclusion, significant differences were observed in the MPV, PDW, P-LCR, PCT and PLT in patients with dengue infection. Low platelet count, low PCT and high PDW may be used as predictor of severity of Dengue infection. Low MPV  $< 9$  fl, low PCT  $< 0.15\%$ , high P-LCR  $> 42\%$  and high PDW  $> 15$  fl shows considerable sensitivity and specificity for dengue fever.

**Keyword:** Mean platelet volume, Dengue, thrombocytopenia, Platelet distribution width, Bikaner

## Introduction

Dengue fever is an acute infectious disease caused by four serotypes of dengue virus, and is the most prevalent mosquito-borne viral disease in humans, occurring in tropical and subtropical countries of the world where over 2.5 billion people are at risk of infection.<sup>1</sup> The World Health Organization has estimated 50 million cases of dengue fever and several hundred thousand cases of dengue hemorrhagic fever occur each year, depending on the epidemic activity.<sup>2</sup> Some 1.8 billion of the population at risk for dengue worldwide live in member states of the WHO South-East Asia Region and Western Pacific Region, which bear nearly 75% of the current global disease burden due to dengue.<sup>3</sup>

Dengue has a wide spectrum of clinical presentations, often with unpredictable clinical evolution and outcome. While most patients recover following a self-limiting non severe clinical course, a small proportion progress to severe disease, mostly characterized by plasma leakage with or without hemorrhage. Early recognition of dengue is challenging because the initial symptoms are often non-specific, viremia may be below detectable levels and serological tests confirm dengue late in the course of illness.<sup>4</sup> The World Health Organization (WHO) classified the clinical presentations of DHF into four severity grades based on laboratory data: Grade I: fever with positive tourniquet test; Grade II: plus mild spontaneous bleeding; Grade III: presence of weak and rapid pulse; and Grade IV: profound shock with undetectable pulse. The last two are considered DSS.<sup>5</sup> Prompt diagnosis during the febrile stage is essential for adjusting appropriate management.<sup>6</sup>

Severity of the illness is determined by various risk factors such as age, pre-existing illness, infecting serotype, and secondary infection. A second infection with a different serotype leads to more severe form of the disease than the primary infection.<sup>7</sup> One of the most common laboratory findings in dengue is thrombocytopenia. The complex mechanism of thrombocytopenia remains unclear. Possible mechanisms of thrombocytopenia could be, direct bone marrow suppression by the virus; anti-dengue antibody-mediated platelet destruction, peripheral consumption of platelets and isolated viral replication in the platelet. Thrombocytopenia leads to bleeding although the platelet count may not directly correlate with the bleeding manifestation.<sup>8</sup>

Recently, novel platelet indices such as MPV, PDW, and P-LCR have been investigated as prospective platelet activation markers.<sup>9</sup> Platelet volume, a marker of platelet function and activity is measured as mean platelet volume (MPV) by hematology analyzers. MPV can be used as independent predictors of bleeding. It is surrogate marker of bone marrow activity; a high MPV indicates increased megakaryocytic activity. A low MPV indicates marrow suppression and increased risk of bleeding. Correlation of platelet count and MPV with bleeding and severity of the disease can potentially predict outcome.<sup>10</sup> Platelets with increased number and size of pseudopodia differ in size, possibly affecting platelet distribution width (PDW) which increases during platelet activation.<sup>9</sup> Furthermore, platelet activation alters the morphology of these cells, which can be evaluated on the basis of mean platelet volume (MPV) and platelet distribution width (PDW), Another platelet parameter is plateletcrit (PCT), which is a reliable measurement of platelet biomass because it combines the MPV with the absolute platelet count.<sup>11</sup> Platelet large cell ratio (P-LCR) was significantly decreased in patients with thrombocytosis than in normal while it was increased in thrombocytopenia. P-LCR was inversely related to platelet count and directly related to PDW and

MPV.<sup>12</sup> The aim of our study was to investigate the platelet indices in patients compared to the control and to assess the role of it the severity in dengue infection and its diagnostic significance.

### **Material and Method**

This study was conducted prospectively for a period from July 2015 to November 2015 during the recent outbreak of dengue in Sardar Patel Medical college, Bikaner, Rajasthan, India. The platelet parameters were measured by the Sysmex KX-21N (Sysmex Corporation, B 7151, MF 9/2008, Japan) semi automated hematology analyzer on venous samples collected in EDTA from 200 patients as a clinical sample and 200 apparent healthy normal individuals as control sample. The inclusion criteria were all patients with clinical features and serologically positive dengue infection included. The exclusion criteria include patient's serologically negative dengue and if routine laboratory testing suggested a bacterial, parasite or any viral infection other than dengue infection or any other disease. The specimens were analysed within 1 hour from venesection. The parameters analysed included PLT, MPV, PDW, P-LCR and PCT. Observations were considered valid only if the specimens were analysed within 1 hour from venesection, to avoid the problems occurring when EDTA collected samples are analysed.

Patients were classified as dengue fever, dengue hemorrhagic fever or dengue shock syndrome according to WHO guidelines and laboratory diagnosis of dengue was established by demonstration of IgM and IgG immune chromatographic Rapid strip test (Pan Bio, Australia) sensitivity 95.6 and specificity 96.

#### *Statistical analysis*

Data were analyzed using a statistical software GraphPad Prism version 6.05 for window. Measurements of laboratory data platelet parameters of patients with DF, DHF were statistically tested by compare mean and Chi-square test which ever was appropriate. A P value less than 0.05 were considered statistically significant. The sensitivity and specificity of relevant platelet indices parameters as an indicator of dengue infection were also assessed by performing receptor operating curve (ROC) analysis.

#### *Ethical Consideration*

Ethical clearance of this study was approved from the regional Ethical Review Committee (ERC). Data regarding the age and sex was recorded in predesigned forms.

### **Result**

This is a case control analytical study conducted in Sardar Patel Medical college, Bikaner, Rajasthan, India. The total number of the confirmed diagnosed dengue patients were 200. The age of the patients in this study was between 16 – 65 years (mean age  $29.69 \pm 12.87$  years). 200 individuals, age and sex matched, were selected as control group. The control individual aged between 16 – 65 years (mean age  $40.43 \pm 16.22$  years). Of the 200 clinical patients, (132) 66% were males and (68) 34% were female. In control group, (120) 60% were males and (80) 40% were females. Demographics data were obtained from patients with dengue virus infection include residence and occupation. 74% patients belongs from Urban area and 26% belong from Rural area. The student was the most common segment of occupation affected (39%), followed by house wife (18), the laborer (17%) and traders (8%).

All of the Patients presented were analysed for PLT, PCT, MPV, PDW, and P-LCR. Median MPV was 8.25 (range 5.7 – 10.4 fl), median PDW was 16.8 (range 14.2 – 18.9 fl), median P-LCR was 46.18 (range 16.1 – 49.1%), median PLT was  $36 \times 10^3/\mu\text{l}$  (range  $12 \times 10^3/\mu\text{l}$  –  $196 \times 10^3/\mu\text{l}$ ) and median PCT was 0.038% (0.012-0.167). Low MPV which indicates bone marrow suppression was noted in 71.95% of patients with DF and 61.11% in patients with DHF. A high PDW which indicates as useful marker for platelet activation was seen in 73.17% of patients with DF and 100% in DHF patients. High P-LCR that correlated with thrombocytopenia was observed in 84.15% of patients with DF and 77.78% in DHF patients (Table 1).

The differences between the patient group and the control group in MPV, PCT and PLT were found to be significantly lower in the patient group ( $P < 0.0001$ ), while the differences between patient group and control in PDW and P-LCR were found to be significant higher in patient group ( $P < 0.0001$ ) (Table 2). There was a significantly lower Platelet count and PCT in DHF group as compared to DF group ( $P < 0.0016$  and  $< 0.0166$  respectively) and significantly higher PDW in DHF group as compared DF group ( $P < 0.0008$ ) (Table 3). There was a significant negative correlation with mean platelet count and PDW and P-LCR ( $P < 0.0017$ ;  $r = -0.3097$  and  $P < 0.0001$ ;  $r = -0.8005$ ) and significant positive correlation with mean platelet count and PCT ( $P < 0.0001$ ;  $r = 0.6829$ ) (Table 4). Table 5 shows that high P-LCR and low PCT more sensitive for DHF than DF and equally specific for both DF and DHF. High PDW is more sensitive for DHF and more specific for DF and low MPV is more sensitive for DF and equally specific for DF and DHF.

## Discussion

Dengue fever is one of the major public health problems. Defect in DF/DHF are multifactorial mechanisms that include thrombopathy, coagulopathy and vasculopathy. Thrombopathy means the thrombocytopenia and platelet dysfunction. Many factors can contribute to the onset of thrombocytopenia in DF from a reactive immune response against platelets to decreased platelet production. The depression in the bone marrow observed in dengue fever in the acute stage may account for thrombocytopenia. In addition, direct infection of megakaryocytes by dengue virus could lead to an increased destruction of platelets.<sup>13</sup> Platelets are involved in hemostasis, tissue repairing, and infection. To our knowledge, there are only few studies investigating changes in platelet indices during dengue infection. MPV has been evaluated as a diagnostic tool in different conditions with thrombocytopenia with contradictory results. It has been demonstrated that MPV has sufficient sensitivity and specificity to discriminate aplastic anemia, bone marrow disease, hypoproliferative thrombocytopenia, and bone marrow metastasis of solid tumor.<sup>14-17</sup> However, it has been reported that although MPV may be used as an initial suggestion of bone marrow disease in thrombocytopenic patient, it has limited sensitivity and specificity.<sup>18</sup> As the platelets are natural sources of growth factors like platelet-derived growth factor (PDGF), vascular endothelial growth factor (VEGF), insulin-like growth factor 1 (IGF-1) or transforming growth factor  $\beta$  (TGF- $\beta$ ), they have important role in inflammation, angiogenesis, repair and regeneration of the tissues.<sup>19</sup>

In our study we found that MPV, Plateletcrit and Platelet count were lower in dengue fever patients as compared to controls and PDW and P-LCR levels were higher in dengue fever patients compared to controls. However, the alteration in MPV level in dengue infection was studied before by Wiwanikit V and found that MPV

in patients with dengue hemorrhagic fever was not decreased and appear to be similar to general healthy population.<sup>14</sup> Bashir AB et al also found that MPV and Platelet count were lower in dengue fever patients when compared to controls and PDW levels were higher in dengue fever patients compared to controls.<sup>20</sup> We noticed that MPV level was significantly lower and PDW level was significantly higher in the dengue fever, which reflect that decreasing MPV and increasing PDW levels may predict dengue fever. Other researchers such as Niethammer et al have upheld the diagnostic value of platelet histogram maximum rather than MPV to differentiate between idiopathic thrombocytopenic purpura (ITP) and hypo productive thrombocytopenia<sup>21,22</sup>. Ntiao et al have concluded that MPV and PDW can be safely relied on for a positive diagnosis of ITP. MPV and PDW were superior to P-LCR.<sup>23</sup> A different approach was adopted by Kurata et al who assessed reticulated platelet which are newly released from the marrow by using RNA-binding dyes and flow cytometric analysis, these immature platelets were successful as a discriminating guide to determine the etiology of thrombocytopenia.<sup>24</sup>

We also found that high P-LCR (>42%) and low PCT(<0.15%) more sensitive for DHF (89% and 100%) than DF(98% and 84%) and equally specific for both DF and DHF(95%). High PDW (>15 fl) is more sensitive for DHF (100% v 95%) and more specific for DF (90% v 85%) and low MPV (<9 fl) is more sensitive for DF (71% v 67%) and equally specific for DF and DHF (95%). Bashir AB et al also found that low MPV <9 fl and high PDW >13fl shows considerable sensitivity for dengue fever.<sup>20</sup> Trupti V. Katti et al found that platelet indices showed linear relationship i.e., decrease (MPV < 6 fl, PDW < 10) with decreasing platelet count in most hypo production group and inverse relationship i.e., Increase (MPV > 10 fl, PDW > 16) with decreasing platelet count was observed in most cases of hyper destruction group.<sup>25</sup> Yanxia Gao et al in their study concluded that the rise in MPV, and to a lesser extent an increase in PLCR and PDW, is indicative of a worse prognosis in patients with septic shock.<sup>26</sup> Mohamed AR et al found that thrombocytopenia occurred in (67.5) % of malarial cases and platelet indices in this study showed (90.2) % had low MPV and (81.3) % had low PCT.<sup>27</sup> We also compared platelet indices between DF and DHF group and found that there was a significantly lower Platelet count and PCT in DHF group as compared to DF group and significantly higher PDW in DHF group as compared DF group. We also found a significant negative correlation with mean platelet count and PDW and P-LCR and significant positive correlation with mean platelet count and PCT.

### **Conclusion**

In conclusion, significant differences were observed in the MPV, PDW, P-LCR, PCT and PLT in patients with dengue infection, low platelet count, low MPV, low PCT, high PDW and high P-LCR may be used as probable indicators for dengue in endemic area. Low platelet count, low PCT and high PDW may be used as predictor of severity of Dengue infection Low MPV <9 fl, low PCT <0.15%, high P-LCR >42% and high PDW >15 fl shows considerable sensitivity and specificity for dengue fever.

**Table 1:** Platelet indices in the case group of the study

Diagnosis	Low MPV	Normal MPV	High MPV	Total
DF	118 (71.95%)	46 (28.05%)	0	164 (82%)
DHF	22 (61.11)	14 (38.89)	0	36 (18%)
Total	140 (70%)	60 (30%)	0	200 (100%)
	Low PDW	Normal PDW	High PDW	Total
DF	0	44 (26.83%)	120 (73.17%)	164 (82%)
DHF	0	0	36 (100%)	36 (18%)
Total	0	44 (22%)	156 (78%)	200 (100%)
	Low P-LCR	Normal P-LCR	High P-LCR	Total
DF	6 (3.66%)	20 (12.19%)	138 (84.15%)	164 (82%)
DHF	2 (5.55%)	6 (16.67%)	28 (77.78%)	36 (18%)
Total	8 (4%)	26 (13%)	166 (83%)	200 (100%)

MPV, mean platelet volume; PDW, platelet distribution width, P-LCR: platelet large cell ratio DF, dengue fever; DHF, dengue haemorrhagic fever

**Table 2:** The difference of platelet indices between case and control groups in the study

Parameter	Test group (mean)	Control group (mean)	P-value	Significance
MPV	8.28 ± 1.012	9.92 ± 0.576	< 0.0001	Extremely significant
PDW	16.69 ± 0.858	14.02 ± 1.060	< 0.0001	Extremely significant
P-LCR	43.96 ± 6.566	32.63 ± 7.363	< 0.0001	Extremely significant
Platelet count	43.87 ± 27.47	291.06 ± 80.41	< 0.0001	Extremely significant
PCT	0.0454 ± 0.0249	0.272 ± 0.0725	< 0.0001	Extremely significant

**Table 3:** Comparison of platelet indices between DF and DHF groups


Parameter	DF (mean)	DHF (mean)	P-value	Significance
MPV (fl)	8.313 ± 1.004	8.133 ± 1.064	< 0.4971	Not significant
PDW (fl)	16.56 ± 0.779	17.29 ± 0.964	< 0.0008	Very significant
P-LCR (%)	43.79 ± 7.131	44.72 ± 2.829	< 0.5913	Not significant
Platelet count (10 <sup>3</sup> /mm <sup>3</sup> )	47.86 ± 28.43	25.67 ± 10.68	< 0.0016	Very significant
PCT (%)	0.0482 ± 0.0259	0.0328 ± 0.0148	< 0.0166	Significant

**Table 4:** Correlation between mean platelet count and platelet indices in study group

Correlation with mean Platelet count	'r' Value	P Value	Significance
PDW	-0.3097	0.0017	Very significant
MPV	0.1668	0.0972	Not significant
P-LCR	-0.8005	<0.0001	Extremely significant
PCT	0.6829	<0.0001	Extremely significant

**Table 5:** Statistical analysis of Platelet indices for diagnosis of Dengue fever (DF) and Dengue hemorrhagic fever (DHF)

Parameters	Sensitivity (%)	Specificity (%)	LR
<b>MPV &lt; 9 fl</b>			
DF	71	95	14.15
DHF	67	95	13.33
<b>PDW &gt; 15 fl</b>			
DF	95	90	9.512
DHF	100	85	6.667
<b>P-LCR &gt; 42 %</b>			
DF	84	95	16.83
DHF	89	95	17.78
<b>PCT &lt; 0.15%</b>			
DF	98	95	19.76
DHF	100	95	20.00



**Figure 1:** Receptor operating characteristic (ROC) curve of MPV, P-LCR, PDW and PCT for dengue infection

## References

1. Halstead SB: Dengue. *Lancet* 2007, 370(9599):1644–1652.
2. Mackenzie JS, Gubler DJ, Petersen LR: Emerging flaviviruses: the spread and resurgence of Japanese encephalitis, West Nile and dengue viruses. *Nat Med* 2004, 10(12 Suppl):S98–S109.
3. Edelman R: Dengue vaccines approach the finish line. *Clinical infectious diseases: an official publication of the Infectious. Dis Soc Am* 2007, 45(Suppl 1):S56–S60.
4. Ramos MM, Tomashek KM, Arguello DF, Luxemburger C, Quinones L, Lang J, Munoz-Jordan JL: Early clinical features of dengue infection in Puerto Rico. *Trans R Soc Trop Med Hyg* 2009, 103(9):878–884
5. WHO, Dengue: Guidelines for Diagnosis, Treatment, Prevention and Control, WHO, Geneva, Switzerland, 2009, [http://whqlibdoc.who.int/publications/2009/9789241547871\\_eng.pdf](http://whqlibdoc.who.int/publications/2009/9789241547871_eng.pdf)
6. Ligon BL: Dengue fever and dengue hemorrhagic fever: a review of the history, transmission, treatment, and prevention. *Semin Pediatr Infect Dis* 2005, 16(1):60–65
7. Guzman MG, Halstead SB, Artsob H, Buchy P, Farrar J, Gubler DJ et al. Dengue: a continuing global threat. *Nat Rev Microbiol.* 2010; 8(12 Suppl): S7 – 16
8. Guota E, Dar L, Kapoor G, Broor S. the changing epidemiology of dengue in delhi, India. *Virology*. 2006; 3:92.
9. Vagdatli E, Gounari E, Lazaridou E, Katsibourlia E, Tsikopoulou F, Labrianou I. Platelet distribution width: a simple, practical and specific marker of activation of coagulation. *HIPPOKRATIA.* 2010; 14 (1): 28-32
10. Wiwanikit V. Mean platelet volume in the patients with dengue hemorrhagic fever. *Platelets.* 2004; 15(3): 185.
11. Greisenegger S, Endler G, Hsieh K, Tentschert S, Mannhalter C, Lalouschek W: Is elevated mean platelet volume associated with a worse outcome in patients with acute ischemic cerebrovascular events? *Stroke* 2004, 35:1688–1691.
12. Babu E, Basu D. Platelet large cell ratio in the differential diagnosis of abnormal platelet counts. *Indian J Pathol Microbiol.* 2004; 47 (2):202 – 5
13. Bashir AB, Saeed OK, Mohammed BA, Ageep AK. Thrombocytopenia and bleeding manifestation among patients with Dengue virus Infection in Port sudan, Red Sea State of Sudan. *J.Infect. Dis. Immun.* . 2015 April ; 7(2) : 7-13
14. Wiwanikit V. Mean platelet volume in the patients with dengue hemorrhagic fever. *Platelets.* 2004; 15(3): 185
15. Babu E, Basu D. Platelet large cell ratio in the differential diagnosis of abnormal platelet counts. *Indian J Pathol Microbiol.* 2004; 47 (2):202 –5
16. Kaito K, Otsubo H, Usui N et al. Platelet size deviation width, platelet large cell ratio, and mean platelet volume have sufficient sensitivity and specificity in the diagnosis of immune thrombocytopenia. *Br Haematol.* 2005; 128: 698 – 702
17. Bowles KM, Cooke LJ, Richards EM, Baglin TP. Platelet size has diagnostic value in patients with thrombocytopenia. *Clin Lab Haem.* 2005; 27: 370 – 373.
18. Numbenjapon T et al. A prospective evaluation of normal mean platelet volume in discriminating hyperdestructive thrombocytopenia from hypoproductive thrombocytopenia. *Int J Lab Hematol* 2008; 30: 408-414.

19. Cabar FR, Fettback PB, Pereira P, Zugaib M. Serum markers in the diagnosis of tubal pregnancy. *Clinics*. 2008;63:701–708
20. Bashir AB, Saeed OK, Mohammed BA, Ageep AK. Role of Platelet Indices in Patients with Dengue Infection in Red Sea State, Sudan. *International Journal of Science and Research (IJSR)*. January 2015 ; 4(1) : 1573-1576
21. Chandra H, Chandra S, Rawat A, Verma SK. Role of mean platelet volume as discriminating guide for bone marrow disease in patients with thrombocytopenia. *Int J Lab Haematol*. 2010; 32 (2)498 – 505.
22. Niethammer AG, Forman EN. Use of platelet histogram maximum in evaluating thrombocytopenia. *Am J Haematol*. 1999; 60: 19 –23.
23. Ntaios G, Papadopoulos A, Chatzinikolaou A et al. Increased value of mean platelet volume and platelet size deviation width may provide a safe positive diagnosis of idiopathic thrombocytopenic purpura. *Acta Haematologica*. 2008; 119:173 –177.
24. Kurata Y et al. diagnostic value of tests for reticulated platelets, plasma glycofibrinogen and thrombopoietin levels for discriminating between hyperdestructive and hypoplastic thrombocytopenia. *Am J Clin Pathol*. 2001; 115: 361 – 369.
25. Trupti V. Katti, Sagar C. Mhetre, Chaya Annigeri. How far are the platelet indices mirror image of mechanism of thrombocytopenia-mystery still remains? *Int J Adv Med*. 2014 Nov;1(3):200-205
26. Yanxia Gao et al. The Impact of Various Platelet Indices as Prognostic Markers of Septic Shock. [www.plosone.org](http://www.plosone.org) . August 2014; 9( 8) e103761:1-7
27. Mohamed AR et al. Evaluation of platelet count and platelet indices in patients with acute Malaria in Shendi locality. *Journal of Biomedical and Pharmaceutical Research* 3 (3) 2014, 01-04